

AD-A283 353



AD _____

GRANT NO: DAMD17-93-J-3006

TITLE: MARKERS OF FIBROBLAST GROWTH FACTOR FAMILY-MEDIATED
GROWTH SIGNAL TRANSDUCTION AS DETERMINANTS OF SUCCESSFUL
HORMONAL THERAPY FOR PATIENTS WITH ESTROGEN RECEPTOR
POSITIVE BREAST CANCER

PRINCIPAL INVESTIGATOR: Francis G. Kern, Ph.D.

CONTRACTING ORGANIZATION: Georgetown University
37th and O Streets, NW
Washington, DC 20057

REPORT DATE: January 14, 1994

TYPE OF REPORT: Annual Report

PREPARED FOR: U.S. Army Medical Research, Development,
Acquisition and Logistics Command (Provisional),
Fort Detrick, Frederick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for public release;
distribution unlimited

The views, opinions and/or findings contained in this report are
those of the author(s) and should not be construed as an official
Department of the Army position, policy or decision unless so
designated by other documentation.

94-25755



1288

94 8 15 014

DTIC
ELECTE
AUG 16 1994
S G D

DTIC QUALITY INSPECTED 2

FOREWORD

Opinions, interpretations, conclusions and recommendations are those of the author and are not necessarily endorsed by the US Army.

Where copyrighted material is quoted, permission has been obtained to use such material.

Where material from documents designated for limited distribution is quoted, permission has been obtained to use the material.

Citations of commercial organizations and trade names in this report do not constitute an official Department of Army endorsement or approval of the products or services of these organizations.

In conducting research using animals, the investigator(s) adhered to the "Guide for the Care and Use of Laboratory Animals," prepared by the Committee on Care and Use of Laboratory Animals of the Institute of Laboratory Resources, National Research Council (NIH Publication No. 86-23, Revised 1985).

For the protection of human subjects, the investigator(s) adhered to policies of applicable Federal Law 45 CFR 46.

In conducting research utilizing recombinant DNA technology, the investigator(s) adhered to current guidelines promulgated by the National Institutes of Health.

In the conduct of research utilizing recombinant DNA, the investigator(s) adhered to the NIH Guidelines for Research Involving Recombinant DNA Molecules.

In the conduct of research involving hazardous organisms, the investigator(s) adhered to the CDC-NIH Guide for Biosafety in Microbiological and Biomedical Laboratories.

Accession For	
NTIS CRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	


PI - Signature

1/14/94
Date

January 14, 1994

Annual: 12/14/92 - 12/14/93

**Markers of Fibroblast Growth Factor Family-Mediated Growth Signal
Transduction as Determinants of Successful Hormonal Therapy for
Patients with Estrogen Receptor Positive Breast Cancer**

G-DAMD17-93-J-3006

Francis G. Kern, Ph.D.

Georgetown University
37th and O Streets, NW
Washington DC 20057

U.S. Army Medical Research, Development,
Acquisition and Logistics Command (Provisional)
Fort Detrick, Frederick, Maryland 21702-5012

Approved for public release,
distribution unlimited

Previous work from the laboratory sponsored by this award demonstrated that overexpression of two members of the fibroblast growth factor (FGF) family of cytokines in a estrogen-dependent human breast cancer cell line resulted in the cancer cells acquiring the ability to grow in mice that were treated with the antiestrogen tamoxifen. Other work demonstrated the presence of high levels mRNA for FGFR-3, one of the transmembrane tyrosine kinase growth factor receptors for FGFs, in a subset of human breast tumors. During the course of the period covered by this report, two high titer polyclonal rabbit antibodies were generated against two peptides present in the FGFR-3 protein. Pathological specimens from 181 patients with estrogen receptor positive breast tumors that received tamoxifen therapy were identified. The suitability of the FGFR-3 antibodies for use in immunohistochemical assays with sections from formalin-fixed and paraffin-embedded material from these patients is currently being evaluated. Reverse transcription- polymerase chain reaction (RT-PCR) was used to generate plasmid transcription vectors for FGF-7 and a FGFR-2 splice variant that generates a specific receptor for FGF-7. These vectors were used to prepare probes for RNase protection assays for FGF-7 and its receptor. Although the probe for the receptor did not appear to be suitable for the assay, the presence of FGF-7 RNA was detected in 35 of 36 human breast tumors examined.

breast cancer, antiestrogen resistance, fibroblast growth factors, growth factor

9

17. SECURITY CLASSIFICATION
OF REPORT

Unclassified

18. SECURITY CLASSIFICATION
OF ABSTRACT

Unclassified

19. SECURITY CLASSIFICATION
OF ABSTRACT

Unclassified

20. SECURITY CLASSIFICATION
OF ABSTRACT

Unlimited

Block 1
Name
Address

Block 2

Block 3

Block 4

Block 5

Block 6

Block 7

Block 8

Block 9

Block 10

Block 11

Block 12

Block 13

Block 14

Block 15

Block 16

Block 17

Block 18

Block 19

Block 20

Block 21

Block 22

Block 23

Block 24

Block 25

Block 26

Block 27

Block 28

Block 29

Block 30

Block 31

Block 32

Block 33

Block 34

Block 35

Block 36

Block 37

Block 38

Block 39

Block 40

Block 41

Block 42

Block 43

Block 44

Block 45

Block 46

Block 47

Block 48

Block 49

Block 50

TABLE OF CONTENTS

Report Documentation Page	1
Table of Contents	2
Introduction	3
Body	4
Conclusions	5
References	6
Appendix	7

INTRODUCTION:

A response to tamoxifen therapy is observed in less than 60% of breast cancer patients with recurrent disease that initially presented with estrogen receptor positive tumors. Therefore additional markers are needed that will allow the clinician to better predict that subpopulation of patients most likely to respond to this therapy. Our laboratory has demonstrated that overexpression of two members of the fibroblast growth factor family in an estrogen receptor positive breast cancer cell line confers these cells with the ability to grow tumors in ovariectomized athymic nude mice treated with tamoxifen (1). These results suggest that FGF signal transduction mediated through FGF binding to specific transmembrane tyrosine kinase receptors may be capable of overcoming the estrogen dependence of estrogen receptor positive (ER+) breast cancers for growth. Indicators of FGF signal transduction being operative within an estrogen receptor positive tumor might therefore be capable of providing a sign of the likelihood that tamoxifen therapy will be ineffective. These indicators could include evidence of expression of either FGF ligands or FGF receptors. Nine members of the FGF family of ligands have been identified to date and four transmembrane tyrosine kinase receptors activated by ligand binding have also been described (2-5). We have used sensitive and specific RNase protection assays to examine the range of expression of six of the nine FGF ligands and all four receptors in human breast tumors. With such an assay we have found that FGFR-3 is highly expressed in a subset of tumors (6). One objective of this project included raising antibodies that could be used in an immunohistochemical assay for FGFR-3 expression. Such antibodies would then be used to perform a retrospective study on formalin fixed and paraffin embedded material from patients with ER+ tumors that received tamoxifen treatment and for whom follow up information could be retrieved. This would allow the one to determine if FGFR-3 overexpression correlates with the failure of tamoxifen therapy. A second objective was to develop RNA probes that could be used to examine the range of expression of FGF-7 expression and its receptor in human breast cancers.

BODY:

A. Preparation of polyclonal rabbit antesera against FGFR-3

1. Choice of peptide antigens. Two hydrophilic, extracellular portions of FGFR3 were chosen as potential antigens using the PlotStructure and Helicalwheel programs in the GCG software package, and using the Antigen program in the PCGENE software package. They are (1) peptide-a: RAAEVPGPPEPGQEEQ, corresponding to amino acids 33-48 (15-mer) of FGFR-3 and (2) peptide-b: PQRLQVLNASMEDS, corresponding to amino acids 91-105 (14-mer) of FGFR-3. These two hydrophilic and extracellular portions were also chosen based on their limited homology to corresponding regions in FGFR-1, FGFR-2 and FGFR-4. 50 mg of HPLC purified peptide-a and peptide-b were synthesized by BioServ, Labs (San Jose, CA).

2. Peptide coupling and rabbit immunization. An additional cysteine residue was attached to the carboxy terminus of both peptides during synthesis to facilitate coupling to carrier. 25 mg of peptide-a and peptide-b were conjugated to Keyhole limpet hemocyanin (KLH) using the heterobifunctional reagent, m-Maleimidobenzoyl-N-hydroxysuccinimide (MBS). For each KLH-coupled peptide, two New Zealand white rabbits, female, weight 3 kg, were immunized by subcutaneous injections into

multiple sites on the rabbit's back. The first immunization was performed with 0.5 mg of peptide-KLH and subsequent boosts were performed with 1mg of peptide-KLH. Complete Freund's adjuvant (GIBCO) was used in the first immunization and incomplete Freund's adjuvant was used in later boosts. The ratio of antigen: adjuvant was 1:1 (v/v). The animals received their first boost three weeks after the primary injection and subsequent boosts were continuously carried out at intervals of 14 days. A prebleed was done immediately before the first immunization. Bleeds from the ear vein and artery were carried out 7 days after boosts.

3. Determination of the titer of anti-FGFR3 in immunized rabbit serum. An enzyme linked immuno absorbant assay (ELISA) for the titer of anti-FGFR-3 was developed as follows. 200 ng/well of peptide-a or peptide-b was added to each well of a 96 well of Reacti-Bind™ Maleic Anhydride Activated Polystyrene Plates (PIERCE) and incubated at 37°C overnight. After washing, 2% bovine serum albumin (BSA) in phosphate buffered saline (PBS) was added at room temperature for 1 hour followed by washing with PBS with 0.1% BSA and 0.05% Tween-20 (wash buffer). The rabbit serum was diluted at 1: 6,000; 1:12,000; 1:24,000; 1: 48,000 in 0.5% BSA-PBS and 200 ul of the diluted immunized rabbit serum was added to the well to allow the anti-FGFR-3 peptide antibodies to bind to corresponding peptide. After incubation at room temperature for 1 hour, the nonspecific protein in the serum was removed by three washes with wash buffer. 200 ul of 1:2000 diluted horseradish peroxidase (HRP) conjugated sheep anti-rabbit IgG was added to the wells and further incubated for 1 hour at room temperature. After three washes to remove unbound second antibodies, 200 ul of HRP substrate, 3,3',5,5' Tetramethyl Benzidine (TMB, 0.4g/L, PIERCE) was added to the well to develop the color. The reaction was stopped with 100 ul of 2M H₂SO₄. The plates was read at A₄₅₀ in Auto-ELISA reader. The prebleed serum at the same dilution was used as background control. The results of the ELISA assays of anti-FGFR-3 from 4 rabbits is included in Figure 1 and Figure 2 in the Appendix. They indicate that for all four sera, positive titers were still obtained at the highest dilution tested (1:512,000).

4. Development of an immunohistochemical assay for FGFR-3.

Information contained within the Georgetown University Medical Center tumor registry was used to generate a list of 181 patients that received tamoxifen therapy. Pathological specimens from approximately 160 of these patients are available through the Georgetown University Medical Center Pathology Department. We are currently in the process of identifying the paraffin blocks from each of these patients that contains breast tumor tissue. Once identified, microscopic sections from each block will be prepared.

B. Development of RNase protection assays for FGF-7 (a.k.a. keratinocyte growth factor or KGF) and the KGF receptor (KGFR) isoform

1. FGF-7. A 195 bp partial cDNA starting from the start codon covering the first 65 amino acids of FGF-7 was synthesized using reverse transcription-polymerase chain reaction (RT-PCR). This portion was chosen based on its relative lack of homology within this region to corresponding regions of FGFs 1-6. The PCR primers utilized were sense : 5'-ATCCACAAATGGATACTGAC-3' and antisense: 5'-TTATATCCCCTCCTTCCATG-3. The template for RT-PCR was human placental RNA. The reverse transcription and PCR reactions were performed according to the

instructions provided by the manufacturer of the RT-PCR kit (Perkin-Elmer). The PCR product was made blunt-ended and ligated into the pGEM7zf+ transcription vector that was linearized with SmaI. the sequence of plasmid inserts from ampicillin resistant colonies arising after transformation was verified by dideoxy sequencing using the T7 P/P primer (Promega). A plasmid having the insert in the orientation: 5'-T7---KGF sense (1-195 bp)---SP6-3' was chosen and plasmid DNA was linearized with EcoRI. A 283 nucleotide P-32 labeled antisense transcript was prepared and RNase protection assays were performed as described in (1). RNA was isolated from a number of human breast cancer cell lines and 36 human breast tumor tissues using either guanidium isothiocyanate or RNazol methods. RNA from the human WI-38 Embryonic lung fibroblast cell line obtained from the ATCC was used as a positive control. All human breast cancer cell lines were negative for expression of FGF-7 mRNA. However, 35 of 36 breast tumor tissues tested were positive in the assay. A photocopy of a representative experiment with breast tumor RNA is included as Figure 3 in the Appendix.

2. Generation of a riboprobe for the FGFR-2 exon IIIc splice variant. Exon shuffling within the third IgG-like domain of FGFR-2 generates a receptor isoform (KGFR) that is capable of binding FGF-7 (KGF) with much higher affinity (5,7). RT-PCR was also used to generate a 171 bp cDNA corresponding to nucleotides 927 to 1098 of the FGFR-2 exon IIIc cDNA, the region that that contains this exon. The primers used were: sense : 5'-CAAGGTTCTCAAGCACTCGGGGATAAATAGTTCCAA-3', and antisense: 5'-CTTCCAGGCGCTTGCTGTTTTGGCAGG-3'. RNA from the MCF-7 ER+ breast carcinoma cell line was used as the template for the reverse transcription. The conditions for PCR were 94°C, 1 min 60°C, 2 min; 72°C, 1 min; 35 cycles. The PCR product was blunt-ended and ligated to SmaI digested pGEM7zf+ DNA and insert sequences were confirmed by dideoxysequencing. A plasmid with the orientation: 5'-T7---KGFR antisense ---SP6-3' was chosen for use. After linearization with Hind III, a 251 nucleotide riboprobe was generated using T7 polymerase. However conditions for RNase protection could not be successfully established with this probe and an artefactual band was consistently observed.

CONCLUSIONS:

High titer FGFR-3 rabbit antisera have successfully been generated and a suitable number of patient samples have been identified. For immunohistochemical assay development we will need positive and negative controls. For this purpose we will use human breast tumors grown in athymic nude mice. The MDA-MB-231 cell line expresses very low levels of RNA for FGFR-3 (8) and therefore the amount of protein produced should be below the detection limits of the assay. Sections from formalin fixed and paraffin embedded MDA MB231 tumors can therefore be used as negative controls. We are currently in the process of constructing a eukaryotic expression vector containing the full length FGFR-3 cDNA. This vector will be used to generate tumorigenic cell lines that overexpress FGFR-3 for use as a positive control.

A successful RNase protection assay for FGF-7 mRNA expression has been developed. The detection of FGF-7 mRNA in human breast tumor tissues but not in any human breast cancer cell lines implies that the expression emanates from the stromal component of the tumor. The probe for KGFR expression was not found to be useful. An RT-PCR assay for KGFR expression in human

tumors is currently being developed as an alternative.

REFERENCES:

1. McLeskey, S.W., Kurebayashi, J., Honig, S.F., Zwiebel, J., Lippman, M.E., Dickson, R.B., and Kern, F.G. Fibroblast growth factor 4 transfection of MCF-7 cells produces cell lines that are tumorigenic and metastatic in ovariectomized or tamoxifen-treated athymic nude mice. *Cancer Res.*, 53:2168-77., 1993.
2. Baird, A. and Klagsbrun, M. The Fibroblast Growth Factor Family. *Cancer Cells*, 3: 239-243, 1991.
3. Tanaka, A., Miyamoto, K., Minamino, N., Takeda, M., Sato, B., Matsuo, H., and Matsumoto, K. Cloning and characterization of an androgen-induced growth factor essential for the androgen-dependent growth of mouse mammary carcinoma cells. *Proc. Natl. Acad. Sci. U. S. A.*, 89:8928-32., 1992.
4. Miyamoto, M., Naruo, K., Seko, C., Matsumoto, S., Kondo, T., and Kurokawa, T. Molecular cloning of a novel cytokine cDNA encoding the ninth member of the fibroblast growth factor family, which has a unique secretion property. *Mol. Cell Biol.*, 13:4251-9., 1993.
5. Johnson, D.E. and Williams, L.T. Structural and functional diversity in the FGF receptor multigene family. *Adv. Cancer Res.*, 60: 1-41, 1993.
6. Ding, I.Y.F., McLeskey, S.W., Chang, K., Fu, Y.M., Acol, J.C., Shou, M.T., Alitalo, K., and Kern, F.G. Expression of fibroblast growth factors (FGFs) and receptors (FGFRs) in human breast carcinomas. *Proc. Am. Assoc. Cancer Res.*, 33: 269, 1992.(Abstract)
7. Miki, T., Fleming, T.P., Bottaro, D.P., Rubin, J.S., Ron, D., and Aaronson, S.A. Expression cDNA cloning of the KGF receptor by creation of a transforming autocrine loop. *Science*, 251: 72-75, 1991.
8. McLeskey, S.W., Ding, I.Y.F., Lippman, M.E., and Kern, F.G. MDA-MB-134 Breast Carcinoma Cells Overexpress FGF Receptors and Are Growth-Inhibited By FGF Ligands. *Cancer Research*, 54: 523-530, 1994.

Titer of rabbit anti-FGFR3-peptide serum




Figure 1




Figure 2




Figure 3. RNAse protection assay demonstrating expression of FGF-7 (KGF) mRNA in Human Breast Tumors. A riboprobe for glyceraldehyde phosphate dehydrogenase (GAPDH) was included in the hybridization cocktail to control for differences in loading.